

# My Asthma Plan

## 1 My usual asthma medicines

- My preventer inhaler is called \_\_\_\_\_  
\_\_\_\_\_ and its colour is \_\_\_\_\_
- I take \_\_\_\_\_ puff/s of my preventer inhaler in the morning and \_\_\_\_\_ puff/s at night. I do this every day even if I feel well.
- Other asthma medicines I take every day:  
\_\_\_\_\_  
\_\_\_\_\_
- My reliever inhaler is called \_\_\_\_\_  
\_\_\_\_\_ and its colour is \_\_\_\_\_  
I take \_\_\_\_\_ puff/s of my reliever inhaler when I wheeze or cough, my chest hurts or it's hard to breathe.
- My best peak flow is \_\_\_\_\_

If I need my blue inhaler to do any sport or activity, I need to see my doctor or asthma nurse.



## 2 My asthma is getting worse if...


- I wheeze or cough, my chest hurts or it's hard to breathe, **or**
- I need my reliever inhaler (usually blue) three or more times a week, **or**
- My peak flow is less than \_\_\_\_\_, **or**
- I'm waking up at night because of my asthma (this is an important sign and I will book a next day appointment)

### If my asthma gets worse, I will:

- Take my preventer medicines as normal
- And also take \_\_\_\_\_ puff/s of my blue reliever inhaler every four hours
- See my doctor or nurse urgently if I don't feel better within 24 hours



**URGENT!** "If your blue reliever inhaler isn't lasting for four hours you are having an asthma attack and you need to take **emergency action now (see section 3)**"



Remember to use my spacer with my inhaler if I have one.

(If I don't have one, I'll check with my doctor or nurse if it would help me)

### Other things to do if my asthma is getting worse

---



---



---



---



---

## 3 I'm having an asthma attack if...

- My reliever inhaler isn't helping or I need it more than every four hours, **or**
- I can't talk, walk or eat easily, **or**
- I'm finding it hard to breathe, **or**
- I'm coughing or wheezing a lot or my chest is tight/hurts, **or**
- My peak flow is less than \_\_\_\_\_

### If I have an asthma attack, I will:



**Call for help**



**Sit up** — don't lie down. Try to be calm.



Take one puff of my reliever inhaler (with my spacer if I have it) **every 30 to 60 seconds** up to a total of 10 puffs.



**If I don't have my blue inhaler, or it's not helping,** I need to call **999** straightaway.



While I wait for an ambulance I can use my blue reliever again, every 30 to 60 seconds (up to 10 puffs) if I need to.

**Even if I start to feel better,** I don't want this to happen again, so I need to see my doctor or asthma nurse **today**.



# My Asthma Plan

## My asthma triggers:

List the things that make your asthma worse and what you can do to help

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

I will see my doctor or asthma nurse at least once a year (but more if I need to)

Date I got my asthma plan:

---

Date of my next asthma review:

---

Doctor/asthma nurse contact details:


---



---

Your asthma plan tells you what medicines to take to stay well

And what to do when your asthma gets worse




## Parents – get the most from your child’s action plan

- **Take a photo** and keep it on your mobile (and your child’s mobile if they have one)
- **Stick a copy** on your fridge door
- **Share** your child’s action plan with school

Learn more about what to do during an asthma attack  
[www.asthma.org.uk/advice/asthma-attacks](http://www.asthma.org.uk/advice/asthma-attacks)

Always keep your reliever inhaler (usually blue) and your spacer with you.

You might need them if your asthma gets worse.



## Questions? Ask Asthma UK’s nurses:

Call on

0300 222 5800 (9am–5pm; Mon–Fri)

Or message on WhatsApp

07378 606 728 (9am–5pm; Mon–Fri)

Name: